arXiv:1507.05241v1 [math.GT] 19 Jul 2015

NONABELIAN REPRESENTATIONS AND SIGNATURES OF DOUBLE TWIST KNOTS

ANH T. TRAN

Dedicated to Professor Jozef Przytycki on the occasion of his 60th birthday

Abstract. A conjecture of Riley about the relationship between real parabolic representations and signatures of two-bridge knots is verified for double twist knots.

1. Introduction

The motivation of this note is a conjecture of Riley about the relationship between real parabolic representations and signatures of two-bridge knots. Two-bridge knots are those knots admitting a projection with only two mixama and two minima. The double branched cover of S^3 along a two-bridge knot is a lens space $L(p, q)$, which is obtained by doing a p/q surgery on the unknot. Such a two-bridge knot is denoted by $\mathfrak{b}(p, q)$. Here p and q are relatively prime integers, and one can always assume that $p > q \geq 1$. It is known that $\mathfrak{b}(p', q')$ is ambient isotopic to $\mathfrak{b}(p, q)$ if and only if $p' = p$ and $q' \equiv q^{\pm 1} \pmod{p}$, see e.g. [BZ]. The knot group of the two-bridge knot $\mathfrak{b}(p, q)$ has a presentation of the form $\langle a, b \mid wa = bw \rangle$ where a, b are meridians, $w = a^{\varepsilon_1} b^{\varepsilon_2} \cdots a^{\varepsilon_{p-2}} b^{\varepsilon_{p-1}}$ and $\varepsilon_j = (-1)^{\lfloor jq/p \rfloor}$. We will call this presentation the Schubert presentation of the knot group of $\mathfrak{b}(p, q)$.

We now study representations of knot groups into $SL_2(\mathbb{C})$. A representation is called nonabelian if its image is a nonabelian subgroup of $SL_2(\mathbb{C})$. Suppose $\rho : \pi_1(\mathfrak{b}(p, q)) \to SL_2(\mathbb{C})$ is a nonabelian representation. Up to conjugation, one can assume that

$$\rho(a) = \begin{bmatrix} s & 1 \\ 0 & s^{-1} \end{bmatrix} \quad \text{and} \quad \rho(b) = \begin{bmatrix} s & 0 \\ 2-y & s^{-1} \end{bmatrix}$$

where $s \neq 0$ and $y \neq 2$ satisfy the matrix equation $\rho(wa) = \rho(bw)$. Riley [Ri1] showed that this matrix equation is actually equivalent to a single polynomial equation $\widetilde{\Phi}_{\mathfrak{b}(p,q)}(s, y) = 0$. The polynomial $\widetilde{\Phi}_{\mathfrak{b}(p,q)}(s, y)$ is called the Riley polynomial of $\mathfrak{b}(p, q)$. It can be chosen so that $\widetilde{\Phi}_{\mathfrak{b}(p,q)}(s, y) = \widetilde{\Phi}_{\mathfrak{b}(p,q)}(s^{-1}, y)$, and hence it can be considered as a polynomial in the variables $x = s + s^{-1}$ and y. Note that $x = \operatorname{tr} \rho(a) = \operatorname{tr} \rho(b)$ and $y = \operatorname{tr} \rho(ab^{-1})$. With these new variables, we write $\Phi_{\mathfrak{b}(p,q)}(x, y)$ for the Riley polynomial of $\mathfrak{b}(p, q)$. See [Le] for an alternative description of $\Phi_{\mathfrak{b}(p,q)}(x, y)$ in terms of traces.

A nonabelian representation of a knot group into $SL_2(\mathbb{C})$ is called parabolic if the trace of a meridian is equal to 2. The zero set in $\mathbb{C}$ of the polynomial $\Phi_K(2, y)$ of a two-bridge knot K describes the set of all parabolic representations of the knot group of K into $SL_2(\mathbb{C})$, see [Ri2]. Riley made the following conjecture about the relationship between the number of real roots of $\Phi_K(2, y)$ and the signature $\sigma(K)$ of K.

2010 *Mathematics Classification:* Primary 57N10. Secondary 57M25.

Key words and phrases: nonabelian representation, parabolic representation, signature, double twist knot.

Conjecture 1. [Ri1] *Suppose K is a two-bridge knot. Then the equation $\Phi_K(2,y)=0$ has at least $\frac{1}{2}|\sigma(K)|$ real solutions.*

In this note we will verify Conjecture 1 for a large class of two-bridge knots, called double twist knots. Let $J(k,l)$ be the two-bridge knot/link in Figure 1, where $k,l \neq 0$ denote the numbers of half twists in the boxes. Positive (resp. negative) numbers correspond to right-handed (resp. left-handed) twists. Note that $J(k,l)$ is a knot if and only if kl is even. The knot $J(2,2n)$ is known as a twist knot. In general, the knot/link $J(k,l)$ is called a double twist knot/link. It is known that

$$\begin{aligned} J(2m,2n) &= \mathfrak{b}(4mn-1,4mn-2n-1),\\ J(2m,-2n) &= \mathfrak{b}(4mn+1,4mn-2n+1),\\ J(2m+1,2n) &= \mathfrak{b}(4mn+2n-1,4mn-1),\\ J(2m+1,-2n) &= \mathfrak{b}(4mn+2n+1,4mn+1) \end{aligned}$$

for positive integers m and n. Moreover, their signatures are 2, 0, $2-2n$ and $2n$ respectively, see e.g. [Mu]. For more information about $J(k,l)$, see [HS, MPL].



FIGURE 1. The two-bridge knot/link $J(k,l)$.

The main results of this note are as follows.

Theorem 1. *Suppose m and n are positive integers. If $x_0 \in \mathbb{R}$ satisfies $\sqrt{4-(mn)^{-1}} < |x_0| \le 2$, then*
(i) the equation $\Phi_{J(2m,2n)}(x_0,y)=0$ has exactly one real solution y, and
(ii) the equation $\Phi_{J(2m,-2n)}(x_0,y)=0$ has no real solutions y,

Theorem 2. *Suppose m and n are positive integers. If $x_0 \in \mathbb{R}$ satisfies $|x_0| \ge 2\cos\frac{\pi}{4m+2}$, then*
(i) the equation $\Phi_{J(2m+1,2n)}(x_0,y)=0$ has at least $n-1$ real solutions y, and
(ii) the equation $\Phi_{J(2m+1,-2n)}(x_0,y)=0$ has at least n real solutions y.

As a consequence of Theorems 1 and 2 we have the following.

Corollary 1. *Conjecture 1 holds true for double twist knots.*

Results in Theorems 1 and 2 are related to the problem of determining the existence of real parabolic representations in the study of left-orderability of the fundamental groups of cyclic branched covers of two-bridge knots, see [Hu, Tr]. A group G is called left-orderable if there exists a strict total ordering $<$ on its elements such that $g<h$ implies $fg<fh$ for all elements $f,g,h \in G$. Left-orderable groups have attracted much attention because of its connection to L-spaces, a class of rational homology 3-spheres defined by Ozsvath and Szabo [OS] using Heegaard Floer homology, via a conjecture of Boyer, Gordon and

Watson [BGW]. This conjecture states that an irreducible rational homology 3-sphere is an L-space if and only if its fundamental group is not left-orderable.

In [BGW, Hu] a sufficient condition for the fundamental group of the n^{th} cyclic branched cover of S^3 along a prime knot to be left-orderable is given, in terms of representations of the knot group. In particular, from the proof of [Hu, Theorem 4.3] it follows that if the knot group of a two-bridge knot K has a real parabolic representation then the n^{th} cyclic branched cover of S^3 along K has left-orderable fundamental group for sufficiently large n. Consequently, if Conjecture 1 holds true then we would have the following: *if K is a two-bridge knot with nonzero signature then the n^{th} cyclic branched cover of S^3 along K has left-orderable fundamental group for sufficiently large n.*

The rest of this note is devoted to the proofs of Theorems 1 and 2. In each proof we begin by computing the knot group and then compute the Riley polynomial $\Phi_K(x,y)$ of a double twist knot K. Finally we study real roots of the polynomial $\Phi_K(2,y)$.

2. Proof of Theorem 1

2.1. $J(2m, 2n)$.

The knot $J(2m,2n)$ is the two-bridge knot $\mathfrak{b}(4mn-1, 4mn-2n-1)$.

2.1.1. *The knot group and the Riley polynomial.*

Lemma 2.1. *Suppose* $1 \le j \le 4mn-2$. *Write* $j = 2mq + r$, *where* $0 \le r \le 2m-1$. *Then* $\varepsilon_j = (-1)^{q+r-1}$.

Proof. We have

$$\frac{(4mn-2n-1)j}{4mn-1} = 2mq + r - \frac{2n(2mq+r)}{4mn-1} = (2m-1)q + r - \frac{q+2nr}{4mn-1}.$$

Since $0 < \frac{q+2nr}{4mn-1} < 1$, we obtain $\lfloor \frac{(4mn-2n-1)j}{4mn-1} \rfloor = (2m-1)q + r - 1$. Hence $\varepsilon_j = (-1)^{q+r-1}$. □

Lemma 2.1 implies that the Schubert presentation of the knot group of $J(2m,2n)$ is $\pi_1(J(2m,2n)) = \langle a, b \mid wa = bw \rangle$ where

$$w = a(b^{-1}a)^{m-1}\big[(ba^{-1})^m(b^{-1}a)^m\big]^{n-1}(ba^{-1})^{m-1}b.$$

The Riley polynomial is computed from the matrix equation $\rho(wa) = \rho(bw)$. The proof of the following is similar that of [MT, Proposition 2.5] and hence is omitted.

Proposition 2.2. *One has* $\Phi_{J(2m,2n)}(x,y) = S_n(t) - \mu S_{n-1}(t)$ *where*

$$\begin{aligned} t &= 2 + (y-2)(y+2-x^2)S^2_{m-1}(y), \\ \mu &= 1 + (y+2-x^2)S_{m-1}(y)(S_m(y) - S_{m-1}(y)). \end{aligned}$$

Here $S_k(z)$'s are the Chebychev polynomials of the second kind defined by $S_0(z) = 1$, $S_1(z) = z$ and $S_k(z) = zS_{k-1}(z) - S_{k-2}(z)$ for all integers k. Note that $S_k(2) = k+1$ and $S_k(-2) = (-1)^k(k+1)$. Moreover if $z = \lambda + \lambda^{-1}$, where $\lambda \neq \pm 1$, then $S_k(z) = \frac{\lambda^{k+1} - \lambda^{-(k+1)}}{\lambda - \lambda^{-1}}$.

In the proofs of Theorems 1 and 2 below, we will need the following properties of $S_k(z)$'s whose proofs can be found, for example, in [Tr] and references therein.

(1) $S^2_k(z) + S^2_{k-1}(z) - zS_k(z)S_{k-1}(z) = 1$.

(2) If z is a real number between -2 and 2, then $|S_{k-1}(z)| \le |k|$.

(3) $S_k(z) = \prod_{j=1}^k (z - 2\cos\frac{j\pi}{k+1})$ and $S_k(z) - S_{k-1}(z) = \prod_{j=1}^k (z - 2\cos\frac{(2j-1)\pi}{2k+1})$ for $k \ge 1$.

2.1.2. *Proof of Theorem 1(i).* Suppose $x_0 \in \mathbb{R}$ satisfies $\sqrt{4-(mn)^{-1}} < |x_0| \le 2$. We want to prove that the equation $\Phi_{J(2m,2n)}(x_0, y) = 0$ has exactly one real solution y.

The proof consists of two steps.

Step 1: The equation $\Phi_{J(2m,2n)}(x_0, y) = 0$ *has no real solutions* $y \le 2$.

The proof of Step 1 is similar to that of [Tr, Proposition 2.7]. Suppose $y \le 2$ and $\Phi_{J(2m,2n)}(x_0, y) = 0$. Then $S_n(t) = \mu S_{n-1}(t)$. Since $S_n^2(t) + S_{n-1}^2(t) - tS_n(t)S_{n-1}(t) = 1$ we have $(\mu^2 + 1 - \mu t)S_{n-1}^2(t) = 1$.

By a direct calculation we have $\mu^2 + 1 - \mu t = (y + 2 - x_0^2)S_{m-1}^2(y)(t + 2 - x_0^2)$. Hence

$$(y + 2 - x_0^2)S_{m-1}^2(y)(t + 2 - x_0^2)S_{n-1}^2(t) = 1. \tag{2.1}$$

Since $(y-2)(y + 2 - x_0^2)S_{m-1}^2(y) = t - 2$, equation (2.1) implies that

$$(t-2)(t + 2 - x_0^2)S_{n-1}^2(t) = y - 2 \le 0.$$

Hence $x_0^2 - 2 < t \le 2$. Then $(y-2)(y+2-x_0^2) = (t-2)/S_{m-1}^2(y) \le 0$, which implies that $x_0^2 - 2 < y \le 2$. Since $|y| \le 2$, we have $|S_{m-1}(y)| \le m$. Similarly $|S_{n-1}(t)| \le n$. Hence

$$(y + 2 - x_0^2)S_{m-1}^2(y)(t + 2 - x_0^2)S_{n-1}^2(t) \le (4 - x_0^2)^2 m^2 n^2 < 1$$

which contradicts equation (2.1).

Step 2: The equation $\Phi_{J(2m,2n)}(x_0, y) = 0$ *has exactly one real solution* $y > 2$.

Consider the following real functions on $(2, \infty)$:

$$\begin{aligned} f(y) &= \Phi_{J(2m,2n)}(x_0, y) = S_n(t) - \mu S_{n-1}(t), \\ g(y) &= (y + 2 - x_0^2)S_{m-1}^2(y)(t + 2 - x_0^2)S_{n-1}^2(t) - 1. \end{aligned}$$

Note that $g(y)$ comes from the equation (2.1). In Step 1 we showed that $f(y) = 0$ implies $g(y) = 0$. Since $g(y)$ is an increasing function on $(2, \infty)$, it has at most one root on $(2, \infty)$. Consequently, $f(y)$ has at most one root on $(2, \infty)$. Since

$$\lim_{y \to 2} f(y) = (n+1) - (1 + (4 - x_0^2)m)n = 1 - (4 - x_0^2)mn > 0$$

and $\lim_{y\to\infty} f(y) = -\infty$, we conclude that $f(y)$ has exactly one root on $(2, \infty)$.

Steps 1 and 2 complete the proof of Theorem 1(i).

2.2. $J(2m, -2n)$.

The knot $J(2m, -2n)$ is the two-bridge knot $\mathfrak{b}(4mn+1, 4mn-2n+1)$.

2.2.1. *The knot group and the Riley polynomial.*

Lemma 2.3. *Suppose* $1 \le j \le 4mn$. *Write* $j = 2mq + r$, *where* $0 \le r \le 2m - 1$. *Then* $\varepsilon_j = (-1)^{q+r-1}$ *if* $r \ge 1$, *and* $\varepsilon_j = (-1)^q$ *if* $r = 0$.

Lemma 2.3 implies that the Schubert presentation of the knot group of $J(2m, -2n)$ is $\pi_1(J(2m, -2n)) = \langle a, b \mid wa = bw \rangle$ where

$$w = \left[(ab^{-1})^m (a^{-1}b)^m\right]^n.$$

Proposition 2.4. *One has* $\Phi_{J(2m,-2n)}(x, y) = S_n(t) - \mu S_{n-1}(t)$ *where*

$$\begin{aligned} t &= 2 + (y-2)(y + 2 - x^2)S_{m-1}^2(y), \\ \mu &= 1 - (y + 2 - x^2)S_{m-1}(y)(S_{m-1}(y) - S_{m-2}(y)). \end{aligned}$$

2.2.2. *Proof of Theorem 1(ii).* Suppose $x_0 \in \mathbb{R}$ satisfies $\sqrt{4-(mn)^{-1}} < |x_0| \le 2$. We want to prove that the equation $\Phi_{J(2m,-2n)}(x_0, y) = 0$ has no real solutions y.

The proof consists of two steps.

Step 1: The equation $\Phi_{J(2m,-2n)}(x_0, y) = 0$ has no real solutions $y \le 2$.

The proof of Step 1 is similar to that of $J(2m, 2n)$.

Step 2: The equation $\Phi_{J(2m,-2n)}(x_0, y) = 0$ has no real solutions $y > 2$.

Suppose $y > 2$. Then $t > 2$ and $\mu < 1$. Hence

$$\Phi_{J(2m,-2n)}(x_0, y) = S_n(t) - \mu S_{n-1}(t) > S_n(t) - S_{n-1}(t) > 0.$$

Steps 1 and 2 complete the proof of Theorem 1(ii).

3. Proof of Theorem 2

3.1. $J(2m{+}1, 2n)$**.** The knot $J(2m{+}1, 2n)$ is the two-bridge knot $\mathfrak{b}(4mn{+}2n{-}1, 4mn{-}1)$.

3.1.1. *The knot group and the Riley polynomial.*

Lemma 3.1. *Suppose $1 \le j \le 2n(2m{+}1){-}2$. Write $j = (2m{+}1)q{+}r$, where $0 \le r \le 2m$. Then $\varepsilon_j = (-1)^{r-1}$.*

Lemma 3.1 implies that the Schubert presentation of the knot group of $J(2m+1, 2n)$ is $\pi_1(J(2m+1, 2n)) = \langle a, b \mid wa = bw\rangle$ where

$$w = (ab^{-1})^m \big[(a^{-1}b)^m a^{-1}b^{-1}(ab^{-1})^m\big]^{n-1}(a^{-1}b)^m.$$

Proposition 3.2. *One has $\Phi_{J(2m+1,2n)}(x, y) = S_n(t) - \mu S_{n-1}(t)$ where*

$$\begin{aligned} t &= x^2 - y - (y-2)(y+2-x^2)S_m(y)S_{m-1}(y), \\ \mu &= 1 - (y - x^2 + 2)S_m(y)(S_m(y) - S_{m-1}(y)). \end{aligned}$$

3.1.2. *Proof of Theorem 2(i).* Suppose $x_0 \in \mathbb{R}$ satisfies $|x_0| \ge 2\cos\frac{\pi}{4m+2}$. We want to prove that the equation $\Phi_{J(2m+1,2n)}(x_0, y) = 0$ has at least $n-1$ real solutions y. To achieve this, we will show that there exist real numbers $y_1 < y_2 < \cdots < y_n$ such that $f(y_j)f(y_{j+1}) < 0$ for $1 \le j \le n-1$, where $f(y) = \Phi_{J(2m+1,2n)}(x_0, y)$.

We have $S_n(t) - S_{n-1}(t) = \prod_{j=1}^n (t - 2\cos\frac{(2j-1)\pi}{2n+1})$. Let $t_j = 2\cos\frac{(2j-1)\pi}{2n+1}$. Since

$$S_n(t_j) = \frac{\sin\frac{(2j-1)(n+1)\pi}{2n+1}}{\sin\frac{(2j-1)\pi}{2n+1}} = \frac{\sin\Big(j\pi - \frac{\pi}{2} + \frac{(2j-1)\pi}{2(2n+1)}\Big)}{\sin\frac{(2j-1)\pi}{2n+1}} = (-1)^{j-1}\frac{\cos\frac{(2j-1)\pi}{2(2n+1)}}{\sin\frac{(2j-1)\pi}{2n+1}}$$

we have $(-1)^{j-1}S_n(t_j) > 0$.

Note that $2 - t = (y - x_0^2 + 2)(S_m(y) - S_{m-1}(y))^2$. Choose the largest real solution y_j of $2 - t_j = (y - x_0^2 + 2)(S_m(y) - S_{m-1}(y))^2$.

Lemma 3.3. *One has $x^2 - 2 < y_1 < y_2 < \cdots < y_n$.*

Proof. Let $g(y) = (y - x_0^2 + 2)(S_m(y) - S_{m-1}(y))^2$ where $y \in \mathbb{R}$. We have $g(y_1) = 2 - t_1 < 2 - t_2$. There exists $y > y_1$ such that $g(y) = 2 - t_2$. Hence $y_2 \ge y > y_1$. Other inequalities can be proved in a similar way. □

We have

$$f(y_j) = (1-\mu)S_n(t_j) = (y_j + 2 - x_0^2)S_m(y_j)(S_m(y_j) - S_{m-1}(y_j))S_n(t_j).$$

Since $y_j > x_0^2 - 2 > 2\cos\frac{\pi}{2m+1}$, we have $S_m(y_j) > S_{m-1}(y_j) > 0$. Moreover, $(-1)^{j-1}S_n(t_j) > 0$. Hence $(-1)^{j-1}f(y_j) > 0$. For each $1 \le j \le n-1$, we have

$$f(y_j)f(y_{j+1}) < 0$$

and hence there exists $y'_j \in (y_j, y_{j+1})$ such that $f(y'_j) = 0$. There are at least $n-1$ real solutions y of the equation $f(y) = 0$.

3.2. *$J(2m+1, -2n)$.* The knot $J(2m+1, -2n)$ is the two-bridge knot $\mathfrak{b}(4mn + 2n + 1, 4mn + 1)$.

3.2.1. *The knot group and the Riley polynomial.*

Lemma 3.4. *Suppose $1 \le j \le 2n(2m+1)$. Write $j = (2m+1)q + r$, where $0 \le r \le 2m$. Then $\varepsilon_j = (-1)^{r-1}$ if $r \ge 1$, and $\varepsilon_j = 1$ if $r = 0$.*

Lemma 3.4 implies that the Schubert presentation of the knot group of $J(2m+1, -2n)$ is $\pi_1(J(2m+1, -2n)) = \langle a, b \mid wa = bw \rangle$ where

$$w \;=\; \left[(ab^{-1})^m ab (a^{-1}b)^m\right]^n.$$

Proposition 3.5. *One has $\Phi_{J(2m+1,-2n)}(x, y) = S_n(t) - \mu S_{n-1}(t)$ where*

$$\begin{aligned} t &= x^2 - y - (y-2)(y+2-x^2)S_m(y)S_{m-1}(y), \\ \mu &= 1 + (y+2-x^2)S_{m-1}(y)(S_m(y) - S_{m-1}(y)). \end{aligned}$$

3.2.2. *Proof of Theorem 2(i).* Suppose $x_0 \in \mathbb{R}$ satisfies $|x_0| \ge 2\cos\frac{\pi}{4m+2}$. We want to prove that the equation $\Phi_{J(2m+1,-2n)}(x_0, y) = 0$ has at least n real solutions y. To achieve this, we will show that there exist real numbers $y_0 < y_1 < y_2 < \cdots < y_n$ such that $f(y_j)f(y_{j+1}) < 0$ for $0 \le j \le n-1$, where $f(y) = \Phi_{J(2m+1,-2n)}(x_0, y)$.

Choose t_j and y_j for $1 \le j \le n$ as in the proof of Theorem 2(i). Then $x^2 - 2 < y_1 < y_2 < \cdots < y_n$. We have

$$f(y_j) = (1-\mu)S_n(t_j) = -(y+2-x_0^2)S_{m-1}(y)(S_m(y) - S_{m-1}(y))S_n(t_j),$$

which implies that $(-1)^j f(y_j) > 0$.

Let $y_0 = x_0^2 - 2$. Then $f(y_0) = S_n(2) - S_{n-1}(2) = 1$. For each $0 \le j \le n-1$, we have

$$f(y_j)f(y_{j+1}) < 0$$

and hence there exists $y'_j \in (y_j, y_{j+1})$ such that $f(y'_j) = 0$. There are at least n real solutions y of the equation $f(y) = 0$.

REFERENCES

[BGW] S. Boyer, C. Gordon, and L. Watson, *On L-spaces and left-orderable fundamental groups*, Math. Ann. **356** (2013), no. 4, 1213–1245.

[BZ] G. Burde and H. Zieschang, *Knots*, de Gruyter Stud. Math., vol. 5, de Gruyter, Berlin, 2003.

[HS] J. Hoste and P. Shanahan, *A formula for the A-polynomial of twist knots*, J. Knot Theory Ramifications **13** (2004), no. 2, 193–209.

[Hu] Y. Hu, *Left-orderability and cyclic branched coverings*, Algebr. Geom. Topol. **15** (2015), no. 1, 399–413.

[Le] T. Le, *Varieties of representations and their subvarieties of cohomology jumps for knot groups*, (Russian) Mat. Sb. **184** (1993), 57–82; translation in Russian Acad. Sci. Sb. Math. **78** (1994), 187–209.

[MPL] M. Macasieb, K. Petersen and R. van Luijk, *On character varieties of two-bridge knot groups* Proc. Lond. Math. Soc. (3) **103** (2011), no. 3, 473–507.

- [MT] T. Morifuji and A. Tran, *Twisted Alexander polynomials of 2-bridge knots for parabolic representations*, Pacific J. Math. **269** (2014), no. 2, 433-451.
- [Mu] K. Murasugi, *Classical knot invariants and elementary number theory*, Primes and knots, 167–196, Contemp. Math., 416, Amer. Math. Soc., Providence, RI, 2006.
- [OS] P. Ozsvath and Z. Szabo, *On knot Floer homology and lens space surgeries*, Topology **44** (2005), no. 6, 1281–1300.
- [Ri1] R. Riley, *Nonabelian representations of 2-bridge knot groups*, Quart. J. Math. Oxford Ser. (2) **35** (1984), 191–208.
- [Ri2] R. Riley, *Parabolic representations of knot groups I*, Proc. London Math. Soc. (3) **24** (1972), 217–242.
- [Tr] A. Tran, *On left-orderability and cyclic branched coverings*, J. Math. Soc. Japan **67** (2015), no. 3, 1169–1178.

Department of Mathematical Sciences, University of Texas at Dallas, Richardson, TX 75080, USA

E-mail address: `att140830@utdallas.edu`